

ON CONJUGACY CLASSES OF $\mathrm{SL}(2, q)$

EDITH ADAN-BANTE AND JOHN M. HARRIS

ABSTRACT. Let $\mathrm{SL}(2, q)$ be the group of 2×2 matrices with determinant one over a finite field $\mathcal{F}$ of size q . We prove that if q is even, then the product of any two noncentral conjugacy classes of $\mathrm{SL}(2, q)$ is the union of at least $q - 1$ distinct conjugacy classes of $\mathrm{SL}(2, q)$. On the other hand, if $q > 3$ is odd, then the product of any two noncentral conjugacy classes of $\mathrm{SL}(2, q)$ is the union of at least $\frac{q+3}{2}$ distinct conjugacy classes of $\mathrm{SL}(2, q)$.

1. INTRODUCTION

Let $\mathcal{G}$ be a finite group, $A \in \mathcal{G}$ and $A^{\mathcal{G}} = \{A^B \mid B \in \mathcal{G}\}$ be the conjugacy class of A in $\mathcal{G}$. Let $\mathcal{X}$ be a $\mathcal{G}$ -invariant subset of $\mathcal{G}$, i.e. $\mathcal{X}^A = \{B^A \mid B \in \mathcal{X}\} = \mathcal{X}$ for all $A \in \mathcal{G}$. Then $\mathcal{X}$ can be expressed as a union of n distinct conjugacy classes of $\mathcal{G}$, for some integer $n > 0$. Set $\eta(\mathcal{X}) = n$.

Given any conjugacy classes $A^{\mathcal{G}}, B^{\mathcal{G}}$ in $\mathcal{G}$, we can check that the product $A^{\mathcal{G}}B^{\mathcal{G}} = \{XY \mid X \in A^{\mathcal{G}}, Y \in B^{\mathcal{G}}\}$ is a $\mathcal{G}$ -invariant subset and thus $A^{\mathcal{G}}B^{\mathcal{G}}$ is the union of $\eta(A^{\mathcal{G}}B^{\mathcal{G}})$ distinct conjugacy classes of $\mathcal{G}$.

Fix a prime p and an integer $m > 0$. Let $\mathcal{F} = \mathcal{F}(q)$ be a field with $q = p^m$ elements and $\mathcal{S} = \mathrm{SL}(2, q) = \mathrm{SL}(2, \mathcal{F})$ be the special linear group, i.e. the group of 2×2 invertible matrices over $\mathcal{F}$ with determinant 1.

It is proved in [1] that for any integer $n > 5$, given any nontrivial conjugacy classes α^{S_n} and β^{S_n} of the symmetric group S_n of n letters, that is $\alpha, \beta \in S_n \setminus \{e\}$, if n is a multiple of two or of three, the product $\alpha^{S_n}\beta^{S_n}$ is the union of at least two distinct conjugacy classes, i.e. $\eta(\alpha^{S_n}\beta^{S_n}) \geq 2$, otherwise the product $\alpha^{S_n}\beta^{S_n}$ is the union of at least three distinct conjugacy classes, i.e. $\eta(\alpha^{S_n}\beta^{S_n}) \geq 3$. A similar result is proved for the alternating group A_n in [2].

Fix a prime p and an integer $m > 0$. Let $\mathcal{F} = \mathcal{F}(q)$ be a field with $q = p^m$ elements and $\mathcal{S} = \mathrm{SL}(2, q) = \mathrm{SL}(2, \mathcal{F})$ be the special linear group, i.e. the group of 2×2 invertible matrices over $\mathcal{F}$ with determinant 1. Given any non-central conjugacy classes $A^{\mathcal{S}}, B^{\mathcal{S}}$ in $\mathcal{S}$, is there any relationship between $\eta(A^{\mathcal{S}}B^{\mathcal{S}})$ and q ?

Arad and Herzog conjectured in [3] that the product of two nontrivial conjugacy classes is never a conjugacy class in a finite nonabelian simple group. Thus, when $q \geq 4$ is even we have that $\mathcal{S} = \mathrm{SL}(2, q) = \mathrm{PSL}(2, q)$ is simple and so we must have that $\eta(A^{\mathcal{S}}B^{\mathcal{S}}) > 1$ unless $A = I$ or $B = I$. In what follows, we expand and refine this statement.

Theorem A. *Fix a positive integer m . Let A and B be matrices in $\mathcal{S} = \mathrm{SL}(2, 2^m)$. Then exactly one of the following holds:*

Date: 2009.

1991 Mathematics Subject Classification. 15A33, 20G40, 20E45.

Key words and phrases. conjugacy classes, matrices over a finite field, products of conjugacy classes, special lineal group.

- (i) $A^S B^S = (AB)^S$ and at least one of A, B is a scalar matrix.
- (ii) $A^S B^S$ is the union of at least $2^m - 1$ distinct conjugacy classes, i.e. $\eta(A^S B^S) \geq 2^m - 1$.

Theorem B. Fix an odd prime p and an integer $m > 0$ such that $q = p^m > 3$. Let A and B be matrices in $\mathcal{S} = \text{SL}(2, q)$. Then exactly one of the following holds:

- (i) $A^S B^S = (AB)^S$ and at least one of A, B is a scalar matrix.
- (ii) $A^S B^S$ is the union of at least $\frac{q+3}{2}$ distinct conjugacy classes, i.e. $\eta(A^S B^S) \geq \frac{q+3}{2}$.

Given any group G , denote by $\min(G)$ the smallest integer in the set $\{\eta(a^G b^G) \mid a, b \in G \setminus \mathbf{Z}(G)\}$. In Proposition 2.13, given any integer $m > 0$, we present matrices A, B in $\text{SL}(2, 2^m)$ such that $\eta(A^{\text{SL}(2, 2^m)} B^{\text{SL}(2, 2^m)}) = 2^m - 1$ and thus Theorem A is optimal. Also, given any $q = p^m > 3$, where p is an odd prime and m is a positive integer, in Proposition 2.20 we prove that Theorem B is optimal by presenting matrices where $\min(\text{SL}(2, q))$ is attained. Also, using GAP [5], we can check that $\min(\text{SL}(2, 3)) = 2$ and thus Theorem B cannot apply when $q = 3$.

When q is even, $\text{SL}(2, q) = \text{PSL}(2, q)$ is a simple group of Lie type of characteristic 2. Hence, if we require that both A and B are not involutions in Theorem A, the conclusion of Theorem A follows from Theorem 2 of [6]. We thank Rod Gow for pointing this out to us.

2. PROOFS

Notation. We will denote with uppercase letters the matrices and with lowercase letters the elements in $\mathcal{F}$.

Remark 2.1. We can describe matrix representatives of conjugacy classes in $\mathcal{S} = \text{SL}(2, \mathcal{F})$ by four families or types ([4]):

- (i) $\begin{pmatrix} r & 0 \\ 0 & r \end{pmatrix}$ where $r \in \mathcal{F}$ and $r^2 = 1$,
- (ii) $\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}$ where $r, s \in \mathcal{F}$ and $rs = 1$.
- (iii) $\begin{pmatrix} s & u \\ 0 & s \end{pmatrix}$ where $s \in \mathcal{F}$, $s^2 = 1$ and u is either 1 or a non-square element of $\mathcal{F}$, i.e. $u \in \mathcal{F} \setminus \{x^2 \mid x \in \mathcal{F}\}$.
- (iv) $\begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}$, where $w = r + r^q$ and $1 = r^{1+q}$ for some $r \in \mathcal{E} \setminus \mathcal{F}$, where $\mathcal{E}$ is a quadratic extension of $\mathcal{F}$.

That is, any conjugacy class A^S of $\mathcal{S}$ must contain one of the above matrices.

Remark 2.2. By Lemma 3 of [1], we have that $A^S B^S = B^S A^S$. Thus if we want to prove that given any non-central conjugacy classes A^S and B^S of $\mathcal{S}$, $\eta(A^S B^S) \geq n$ for some integer n , it suffices to prove that the statement holds for each of the six combinations of conjugacy classes containing matrices of type (ii), (iii) and (iv).

Remark 2.3. Two matrices in the same conjugacy class have the same trace. Thus, if the matrices do not have the same trace, then they belong to distinct conjugacy classes.

Lemma 2.4. Let $C = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathcal{S}$ and $A = \begin{pmatrix} e & f \\ g & h \end{pmatrix} \in \mathcal{S}$. Then

$$A^C = C^{-1}AC = \begin{pmatrix} a(de - bg) + c(df - bh) & b(de - bg) + d(df - bh) \\ a(-ce + ag) + c(-cf + ah) & b(-ce + ag) + d(-cf + ah) \end{pmatrix},$$

and therefore

(i)

$$\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}^C = \begin{pmatrix} adr - bcs & bd(r - s) \\ -ac(r - s) & ads - bcr \end{pmatrix}.$$

(ii)

$$\begin{pmatrix} s & u \\ 0 & s \end{pmatrix}^C = \begin{pmatrix} s + ucd & ud^2 \\ -uc^2 & s - ucd \end{pmatrix}.$$

(iii)

$$\begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}^C = \begin{pmatrix} ab + c(d - bw) & b^2 + d^2 - bdw \\ -a^2 - c^2 + acw & -ab + d(-c + aw) \end{pmatrix}.$$

Proof. Observe that $C^{-1} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$.

Hence

$$\begin{aligned} C^{-1}AC &= \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \begin{pmatrix} e & f \\ g & h \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \\ &= \begin{pmatrix} de - bg & df - bh \\ -ce + ag & -cf + ah \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \\ &= \begin{pmatrix} a(de - bg) + c(df - bh) & b(de - bg) + d(df - bh) \\ a(-ce + ag) + c(-cf + ah) & b(-ce + ag) + d(-cf + ah) \end{pmatrix}. \end{aligned}$$

□

Lemma 2.5. Let $C = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathcal{S}$. Then

$$(i) \text{ Trace}\left(\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}^C \begin{pmatrix} u & 0 \\ 0 & v \end{pmatrix}\right) = ad(r - s)(u - v) + (us + vr).$$

$$(ii) \text{ Trace}\left(\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}^C \begin{pmatrix} t & u \\ 0 & t \end{pmatrix}\right) = t(r + s) - ac(r - s)u.$$

$$(iii) \text{ Trace}\left(\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}^C \begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}\right) = (ac + bd)(s - r) + w(ads - bcr).$$

$$(iv) \text{ Trace}\left(\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^C \begin{pmatrix} t & w \\ 0 & t \end{pmatrix}\right) = 2rt - uwc^2.$$

$$(v) \text{ Trace}\left(\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^C \begin{pmatrix} 0 & 1 \\ -1 & s \end{pmatrix}\right) = -ud^2 - uc^2 + s(r - ucd).$$

$$(vi) \text{ Trace}\left(\begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}^C \begin{pmatrix} 0 & 1 \\ -1 & v \end{pmatrix}\right) = -a^2 - b^2 - c^2 - d^2 + bdw + acw + v(-ab + d(-c + aw)).$$

Proof. The result follows by Lemma 2.4 and

(i)

$$\begin{aligned}
\text{Trace}\left(\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}^C \begin{pmatrix} u & 0 \\ 0 & v \end{pmatrix}\right) &= \text{Trace}\left(\begin{pmatrix} adr - bcs & bd(r-s) \\ -ac(r-s) & ads - bcr \end{pmatrix} \begin{pmatrix} u & 0 \\ 0 & v \end{pmatrix}\right) \\
&= u(adr - bcs) + v(ads - bcr) \\
&= ad(ur + vs) - bc(us + vr) \\
&= ad(ur + vs) + (1 - ad)(us + vr) \\
&= ad(ur + vs - us - vr) + (us + vr) \\
&= ad(u(r-s) - v(r-s)) + (us + vr) \\
&= ad(u-v)(r-s) + (us + vr).
\end{aligned}$$

(ii)

$$\begin{aligned}
\text{Trace}\left(\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}^C \begin{pmatrix} t & u \\ 0 & t \end{pmatrix}\right) &= \text{Trace}\left(\begin{pmatrix} adr - bcs & bd(r-s) \\ -ac(r-s) & ads - bcr \end{pmatrix} \begin{pmatrix} t & u \\ 0 & t \end{pmatrix}\right) \\
&= t(adr - bcs) - ac(r-s)u + t(ads - bcr) \\
&= ad(rt + st) - bc(st + rt) - ac(r-s)u \\
&= (ad - bc)t(r+s) - ac(r-s)u \\
&= t(r-s) - ac(r-s)u.
\end{aligned}$$

(iii)

$$\begin{aligned}
\text{Trace}\left(\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}^C \begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}\right) \\
&= \text{Trace}\left(\begin{pmatrix} adr - bcs & bd(r-s) \\ -ac(r-s) & ads - bcr \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}\right) \\
&= -bd(r-s) - ac(r-s) + w(ads - bcr) \\
&= (ac + bd)(s-r) + w(ads - bcr).
\end{aligned}$$

(iv)

$$\begin{aligned}
\text{Trace}\left(\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^C \begin{pmatrix} t & w \\ 0 & t \end{pmatrix}\right) &= \begin{pmatrix} r + ucd & ud^2 \\ -uc^2 & r - ucd \end{pmatrix} \begin{pmatrix} t & w \\ 0 & t \end{pmatrix} \\
&= t(r + ucd) + w(-uc^2) + t(r - ucd) \\
&= 2rt - uwc^2.
\end{aligned}$$

(v)

$$\begin{aligned}
\text{Trace}\left(\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^C \begin{pmatrix} 0 & 1 \\ -1 & s \end{pmatrix}\right) &= \text{Trace}\left(\begin{pmatrix} r + ucd & ud^2 \\ -uc^2 & r - ucd \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & s \end{pmatrix}\right) \\
&= -ud^2 - uc^2 + s(r - ucd)
\end{aligned}$$

(vi)

$$\begin{aligned}
\text{Trace}\left(\begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}^C \begin{pmatrix} 0 & 1 \\ -1 & v \end{pmatrix}\right) \\
&= \text{Trace}\left(\begin{pmatrix} ab + c(d-bw) & b^2 + d^2 - bdw \\ -a^2 - c^2 + acw & -ab + d(-c + aw) \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & v \end{pmatrix}\right) \\
&= -b^2 - d^2 + bdw - a^2 - c^2 + acw + v(-ab + d(-c + aw))
\end{aligned}$$

□

Remark 2.6. For any $a, b \in \mathcal{F}$ such that $a \neq 0$, we have that $\{ax + b \mid x \in \mathcal{F}\} = \mathcal{F}$.

Lemma 2.7. Let $A = \begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}$, where $r \neq s$, and B be any non-central matrix in $\mathcal{S}$, i.e. of type (ii), (iii) or (iv) in Remark 2.1. Given any $f \in \mathcal{F}$, there exists a matrix D in the product $A^{\mathcal{S}}B^{\mathcal{S}}$ such that $\text{Trace}(D) = f$. In particular, $\eta(A^{\mathcal{S}}B^{\mathcal{S}}) \geq q$ for any non-central matrix B .

Proof. Given $i \in \mathcal{F}$, set $C(i) = \begin{pmatrix} i & i-1 \\ 1 & 1 \end{pmatrix}$. Observe that $C(i) \in \mathcal{S}$ for all $i \in \mathcal{F}$. Fix u, v in $\mathcal{F}$ such that $uv = 1$. By Lemma 2.5 (i), we have that

$$(2.8) \quad \text{Trace}\left(\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}^{C(i)} \begin{pmatrix} u & 0 \\ 0 & v \end{pmatrix}\right) = (r-s)(u-v)i + (us+vr).$$

By Lemma 2.5 ii) we have that

$$(2.9) \quad \text{Trace}\left(\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}^{C(i)} \begin{pmatrix} t & u \\ 0 & t \end{pmatrix}\right) = -(r-s)ui + t(r+s).$$

Now let $E(i) = \begin{pmatrix} 1 & i \\ 0 & 1 \end{pmatrix}$ for $i \in \mathcal{F}$. As before, observe that $E(i) \in \mathcal{S}$ for any $i \in \mathcal{F}$. Then by 2.5 (iii), we have that

$$(2.10) \quad \text{Trace}\left(\begin{pmatrix} r & 0 \\ 0 & s \end{pmatrix}^{E(i)} \begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}\right) = (s-r)i + ws.$$

Since $(r-s)(u-v) \neq 0$, $(r-s)u \neq 0$, and $s-r \neq 0$, the result follows from (2.8), (2.9), (2.10), Remark 2.1, Remark 2.3 and Remark 2.6. $\square$

Lemma 2.11. Let $\mathcal{F}$ be a field with 2^m elements for some integer m and $a \in \mathcal{F}$ with $a \neq 0$. Given any $\mathcal{H} \subseteq \mathcal{F}$, the set $\{ai^2 + c \mid i \in \mathcal{H}\}$ has $|H|$ elements. In particular, if $\mathcal{H} = \mathcal{F}$, then the set $\{ai^2 + c \mid i \in \mathcal{H}\}$ has q elements.

Proof. Observe that

$$|\{ai^2 + c \mid i \in \mathcal{H}\}| = |\{ai^2 \mid i \in \mathcal{H}\}| = |\{i^2 \mid i \in \mathcal{H}\}|.$$

Since $\mathcal{F}$ is a field of characteristic 2, the map $x \mapsto x^2$ is an automorphism of $\mathcal{F}$ and thus $|\{i^2 \mid i \in \mathcal{H}\}| = |H|$. $\square$

Lemma 2.12. Let $\mathcal{F}$ be a field with $q = 2^m$ elements for some integer m .

(i) For any f in $\mathcal{F}$, there exists a matrix D in the product $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^{\mathcal{S}} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^{\mathcal{S}}$ such that $\text{Trace}(D) = f$.

(ii) For any f in $\{i^2 + w \mid i \in \mathcal{F} \setminus \{0\}\}$, there exists a matrix D in the product $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^{\mathcal{S}} \begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}^{\mathcal{S}}$ such that $\text{Trace}(D) = f$.

(iii) Given any $f \in \mathcal{F}$, there exists a matrix D in the product $\begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}^{\mathcal{S}} \begin{pmatrix} 0 & 1 \\ -1 & v \end{pmatrix}^{\mathcal{S}}$, where $vw \neq 0$, such that $\text{Trace}(D) = f$.

Thus given any conjugacy classes $A^{\mathcal{S}}, B^{\mathcal{S}}$ in $\mathcal{S}$, where A, B is either of type (iii) or (iv), the product $A^{\mathcal{S}}B^{\mathcal{S}}$ is the union of at least $q-1$ distinct conjugacy classes.

Proof. (i) Given any $i \in \mathcal{F}$, set $C(i) = \begin{pmatrix} 1 & 0 \\ i & 1 \end{pmatrix}$. Observe that $C(i)$ is in $\mathcal{S}$. By Lemma 2.5 (iv)

$$\text{Trace}\left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^{C(i)} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}\right) = i^2.$$

By Lemma 2.11 we have that $\{i^2 \mid i \in \mathcal{F}\} = \mathcal{F}$ and thus (i) follows.

(ii) Given any $i \in \mathcal{F} \setminus \{0\}$, set $E(i) = \begin{pmatrix} i^{-1} & 0 \\ 0 & i \end{pmatrix}$. By Lemma 2.5 (v), we have

$$\text{Trace}\left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^{E(i)} \begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}\right) = i^2 + w.$$

(iii) For $i \in \mathcal{F}$, set $F(i) = \begin{pmatrix} i+1 & i \\ i & i+1 \end{pmatrix}$. Observe that since $\mathcal{F}$ is of characteristic 2, we have $\det(F(i)) = (i+1)^2 - i^2 = i^2 + 1 - i^2 = 1$ and thus $F(i) \in \mathcal{S}$. By Lemma 2.5 (vi) we have

$$\text{Trace}\left(\begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}^{F(i)} \begin{pmatrix} 0 & 1 \\ -1 & v \end{pmatrix}\right) = vw(i^2 + 1).$$

If $vw \neq 0$, then the set $\{vw(i^2 + 1) \mid i \in \mathcal{F}\} = \mathcal{F}$ by Lemma 2.11.

Since $\mathcal{F}$ is of characteristic 2, the only matrix representative of type (iii) is $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Also, since $\mathcal{F}$ is of characteristic two, we can check that the matrices $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ are in the same conjugacy class. Thus, with cases (i), (ii), and (iii), we cover all possible combinations of representatives of type (iii) and (iv), and by Remark 2.3, the proof of the result is complete. $\square$

Proof of Theorem A. If at least one of A or B is in the center, i.e. A or B are of type (i) in Remark 2.1, then $A^{\mathcal{S}}B^{\mathcal{S}} = (AB)^{\mathcal{S}}$. Theorem A then follows from Remark 2.2, Lemma 2.7 and Lemma 2.12. $\square$

Proposition 2.13. *Fix $q = 2^m$ for some integer $m > 0$ and let $\mathcal{F}$ be a field with q elements. Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}$ in $\mathcal{S}$, where $x^2 - wx + 1$ is an irreducible polynomial over $\mathcal{F}$. Then $\eta(A^{\mathcal{S}}B^{\mathcal{S}}) = q - 1$.*

Hence, Theorem A is optimal.

Proof. Set $C = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ in $\mathcal{S}$. By Lemma 2.5 (iv), we have that $\text{Trace}(A^C B) = -d^2 - c^2 + w(1 - cd)$. Since $x^2 - wx + 1$ is an irreducible polynomial over $\mathcal{F}$, $-d^2 - c^2 + w(1 - cd) \neq w$ for any $c, d \in \mathcal{F}$. Thus the matrices in $A^{\mathcal{S}}B^{\mathcal{S}}$ do not have trace w . Also, since the eigenvalues of B are not in $\mathcal{F}$ and the eigenvalues of A is 1, then $A^{\mathcal{S}} \neq (B^{-1})^{\mathcal{S}}$ and so the identity I is not in $A^{\mathcal{S}}B^{\mathcal{S}}$.

Since $\mathcal{F}$ has even characteristic and I is not in $A^{\mathcal{S}}B^{\mathcal{S}}$, we conclude that there is a one-to-one correspondence of the conjugacy classes in $A^{\mathcal{S}}B^{\mathcal{S}}$ with the traces of the matrices: if the trace is 0, then the matrix is similar to a matrix of type (iii), and otherwise, the matrix is similar to a matrix of type (ii) or type (iv), depending on whether or not its characteristic equation is reducible. Thus $\eta(A^{\mathcal{S}}B^{\mathcal{S}}) = q - 1$. $\square$

Lemma 2.14. *Let u, v in $\mathcal{F}$. Then the matrices $\begin{pmatrix} s & v \\ 0 & s \end{pmatrix}$ and $\begin{pmatrix} s & u \\ 0 & s \end{pmatrix}$ are similar if and only if $v = ud^2$ for some d in $\mathcal{F} \setminus \{0\}$. In particular, if u is a*

non-square, then the matrices $\begin{pmatrix} s & 1 \\ 0 & s \end{pmatrix}$ and $\begin{pmatrix} s & u \\ 0 & s \end{pmatrix}$ are not similar, i.e. they belong to distinct conjugacy classes.

Proof. By Lemma 2.4 (ii), we have that $\begin{pmatrix} s & u \\ 0 & s \end{pmatrix}^C = \begin{pmatrix} s + ucd & ud^2 \\ -uc^2 & s - ucd \end{pmatrix}$. Thus, if $\begin{pmatrix} s + ucd & ud^2 \\ -uc^2 & s - ucd \end{pmatrix} = \begin{pmatrix} t & v \\ 0 & t \end{pmatrix}$, then $c = 0$ and $s = t$ or $u = v = 0$ and hence, $v = ud^2$. $\square$

Lemma 2.15. *Let $\mathcal{F}$ be a finite field with q elements and $a, b, c \in \mathcal{F}$ with $a \neq 0$. If q is an odd number, then the set $\{ai^2 + bi + c \mid i \in \mathcal{F}\}$ has exactly $\frac{q+1}{2}$ elements.*

Proof. Since the field $\mathcal{F}$ is of odd characteristic, we have that $2 \neq 0$ and thus

$$\begin{aligned} |\{ai^2 + bi + c \mid i \in \mathcal{F}\}| &= |\{i^2 + \frac{b}{a}i + \frac{c}{a} \mid i \in \mathcal{F}\}| \\ &= |\{i^2 + \frac{b}{a}i + (\frac{b}{2a})^2 \mid i \in \mathcal{F}\}| \\ &= |\{(i + \frac{b}{2a})^2 \mid i \in \mathcal{F}\}| = |\{i^2 \mid i \in \mathcal{F}\}|. \end{aligned}$$

Since q is odd, then 2 divides $q - 1$ and so the square of the set of units forms a subgroup of order $\frac{q-1}{2}$. Since $0^2 = 0$, we have that the set $\{ai^2 + bi + c \mid i \in \mathcal{F}\}$ has exactly $\frac{q-1}{2} + 1 = \frac{q+1}{2}$ elements. $\square$

Lemma 2.16. *Let F be a finite field of size q , where q is odd.*

(i) *Fix $a, b \in \mathcal{F} \setminus \{0\}$, and suppose that $q > 3$. The set $\{ax^2 + by^2 \mid x, y \in \mathcal{F} \setminus \{0\}\}$ has a square and a non-square element.*

(ii) *Fix r, s and u in $\mathcal{F}$ with $s^2 - 4 \neq 0$. Then the set $\{-ux^2 - uy^2 + s(r - uxy) \mid x, y \in \mathcal{F}, (x, y) \neq (0, 0)\}$ has at least $q - 1$ elements.*

(iii) *Let $w \in \mathcal{F}$ be such that $w^2 - 4 \neq 0$. Then $\{a^2 - c^2 + acw \mid a, c \in \mathcal{F}, a \neq 0\} = \mathcal{F} \setminus \{0\}$.*

Proof. (i) Observe that if $c \in \mathcal{F}$ is a non-square element, i.e. $c \notin \{i^2 \mid i \in \mathcal{F}\}$, then $\mathcal{F} = \{i^2 \mid i \in \mathcal{F}\} \cup \{ci^2 \mid i \in \mathcal{F}\}$. Thus, if x is a square and y is a non-square element, then ax is a square and ay is a non-square element when a is a square, and otherwise, ax is a non-square element and ay is a square. Thus the set $\{ax^2 + by^2 \mid x, y \in \mathcal{F} \setminus \{0\}\}$ has a square and a non-square element if and only if $\{x^2 + \frac{b}{a}y^2 \mid x, y \in \mathcal{F} \setminus \{0\}\}$ has a square and a non-square element. Hence, without loss of generality, we may assume that $a = 1$.

Note that there are $\frac{q+1}{2}$ square elements in $\mathcal{F}$ and there are $\frac{q-1}{2}$ non-square elements in $\mathcal{F}$. Also, since $|\mathcal{F}| > 3$, if $\mathcal{F}$ has characteristic $p \neq 3$ then $3^2 + 4^2 = 5^2$, otherwise there exists an element $w \in \mathcal{F}$ such that $w^2 + 1 = 0$. Thus the set $\{x^2 + y^2 \mid x, y \in \mathcal{F}\}$ always contains a square element. If b is a square element, the set $\{x^2 + by^2 \mid x, y \in \mathcal{F} \setminus \{0\}\} = \{x^2 + y^2 \mid x, y \in \mathcal{F}\}$ has a square and a non-square element, otherwise the set of square elements would be a subfield of $\mathcal{F}$ of size $\frac{q+1}{2}$.

Suppose that b is a non-square element. If -1 is a non-square element, then $\{x^2 + by^2 \mid x, y \in \mathcal{F} \setminus \{0\}\} = \{x^2 - y^2 \mid x, y \in \mathcal{F} \setminus \{0\}\}$, and hence we may assume that $b = -1$. Let ϵ be a generator of $\mathcal{F}$. Observe that ϵ is not a square and if $x = \frac{\epsilon+1}{2} = 1 + y$, then $x^2 - y^2 = (x - y)(x + y) = \epsilon$ and $x, y \in \mathcal{F} \setminus \{0\}$ since $|\mathcal{F}| > 3$. Also for any $x = y \neq 0$, we have that $x^2 - y^2 = 0$ and thus the set $\{x^2 - y^2 \mid x, y \in \mathcal{F} \setminus \{0\}\}$ contains a square, namely zero, and a non-square element, namely ϵ . We may assume then that -1 is a square element.

Given $z \in \mathcal{F} \setminus \{0\}$, set $y = xz$. Then $x^2 + by^2 = x^2(1 + bz^2)$. Observe that $1 + bz^2 = 0$ does not have a solution since -1 is square and so $-1/z^2$ is a square. Since the set $Z = \{1 + bz^2 \mid z \in \mathcal{F} \setminus \{0\}\}$ has $\frac{q-1}{2}$ elements and $0, 1 \notin Z$, either Z has a square and a non-square element, or it has only non-square elements. In the first case, since $\{x^2(1 + bz^2) \mid x, z \in \mathcal{F} \setminus \{0\}\} \subset \{x^2 + by^2 \mid x, y \in \mathcal{F}\}$, the result follows. We may assume now that Z is the set of non-square elements. Given $w \in \mathcal{F} \setminus \{0\}$, set $x = wy$. Then $x^2 + by^2 = y^2(w^2 + b)$. Hence $W = \{w^2 + b \mid w \in \mathcal{F} \setminus \{0\}\}$ has $\frac{q-1}{2}$ elements and $b, -b \notin W$. Since both b and $-b$ are non-squares, it follows that either W contains both square and non-square elements, or W contains only square elements. In the first case, the result follows as before. In the second case, since $\{x^2(1 + bz^2) \mid x, z \in \mathcal{F} \setminus \{0\}\}$ is the set of all non-square elements and $\{y^2(w^2 + b) \mid y, w \in \mathcal{F} \setminus \{0\}\}$ is the set of all nonzero square elements, (i) follows.

(ii) Observe that $-ux^2 - uy^2 + s(r - uxy) = f$ if and only if

$$(2.17) \quad x^2 + y^2 - sxy + \frac{f - sr}{u} = 0$$

for some (x, y) . Observe that (2.17) has a solution if there is some $y \in \mathcal{F}$ such that the discriminant $\Delta(y) = (sy)^2 - 4(y^2 + \frac{f-sr}{u}) = (s^2 - 4)y^2 + \frac{4(f-sr)}{u}$ is a square. Since $s^2 - 4 \neq 0$, by Lemma 2.15 we have that the set $\{(s^2 - 4)y^2 + \frac{4(f-sr)}{u} \mid y \in \mathcal{F}\}$ has at least $\frac{q+1}{2}$ elements. Since there are $\frac{q+1}{2}$ squares, for some y we must have that $\Delta(y)$ is a square. It follows then that $x = \frac{sy \pm \sqrt{\Delta(y)}}{2}$ is a solution for (2.17). Observe that $(x, y) = (0, 0)$ is a solution for the equation (2.17) if and only if $f - sr = 0$. We conclude that for at least $q - 1$ elements of $\mathcal{F}$, (2.17) has a solution (x, y) with $(x, y) \neq (0, 0)$.

(iii) Observe that the equation $x^2 + y^2 - xyw = 0$ has the unique solution $(x, y) = (0, 0)$ since $w^2 - 4$ is not a square and so the discriminant $\delta(y) = y^2w^2 - 4y^2 = y^2(w^2 - 4)$ is a square if and only if $y = 0$. As before, we can check that for any $f \in \mathcal{F} \setminus \{0\}$, the equation $x^2 + y^2 - xyw = f$ has a solution with $x \neq 0$. $\square$

Lemma 2.18. *Let $q > 3$ be odd, and let $\mathcal{F}$ be a finite field with q elements.*

(i) *Given any f in $\{2rt - uwi^2 \mid i \in \mathcal{F}\}$, there exists a matrix B in the product $\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^S \begin{pmatrix} t & w \\ 0 & t \end{pmatrix}^S$ such that $\text{Trace}(B) = f$. Also for some $a, b \in \mathcal{F}$, where a*

is a square and b is a non-square, the matrices $\begin{pmatrix} rt & a \\ 0 & rt \end{pmatrix}$ and $\begin{pmatrix} rt & b \\ 0 & rt \end{pmatrix}$ are in the product $\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^S \begin{pmatrix} t & w \\ 0 & t \end{pmatrix}^S$. Thus $\eta(\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^S \begin{pmatrix} t & w \\ 0 & t \end{pmatrix}^S) \geq \frac{q+3}{2}$.

(ii) *For any f in $\{-uc^2 - ud^2 + s(r - cd) \mid c, d \in \mathcal{F} \setminus \{0\}\}$, there exists a matrix D in the product $\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^S \begin{pmatrix} 0 & 1 \\ -1 & s \end{pmatrix}^S$ such that $\text{Trace}(D) = f$. Therefore $\eta(\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^S \begin{pmatrix} 0 & 1 \\ -1 & s \end{pmatrix}^S) \geq q - 1$.*

We conclude that given any conjugacy classes A^S, B^S in $\mathcal{S}$, where at least one of A or B is of type (iii), the product $A^S B^S$ is the union of at least $\frac{q+3}{2}$ distinct conjugacy classes.

Proof. (i) By Lemma 2.4, given any $x, y \in \mathcal{F} \setminus \{0\}$,

$$\begin{pmatrix} rt & rwy^2 + tux^2 \\ 0 & rt \end{pmatrix} = \begin{pmatrix} r & ux^2 \\ 0 & r \end{pmatrix} \begin{pmatrix} t & wy^2 \\ 0 & t \end{pmatrix} \in \begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^S \begin{pmatrix} t & w \\ 0 & t \end{pmatrix}^S$$

By Lemma 2.16 (i), the set $\{rwy^2 + tux^2 \mid x, y \in \mathcal{F} \setminus \{0\}\}$ has a square and a non-square element. It follows then by Lemma 2.14 that there are two matrices that are not similar in the product with the same trace.

Given any $i \in \mathcal{F}$, let $C(i) = \begin{pmatrix} 1 & 0 \\ i & 1 \end{pmatrix}$. By Lemma 2.5 (iv), we have that

$$\text{Trace}\left(\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^{C(i)} \begin{pmatrix} t & w \\ 0 & t \end{pmatrix}\right) = 2rt - uwi^2.$$

By Lemma 2.15, the set $\{2rt - uwi^2 \mid i \in \mathcal{F}\}$ has $\frac{q+1}{2}$ elements. Thus there are at least $\frac{q+1}{2}$ distinct values for the traces of the matrices in the product. Since there are at least two matrices that are not similar in the product with the same trace and $\frac{q+1}{2} + 1 = \frac{q+3}{2}$, (i) follows.

(ii) Let $C = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathcal{S}$. By Lemma 2.5 (v), we have

$$\text{Trace}\left(\begin{pmatrix} r & u \\ 0 & r \end{pmatrix}^C \begin{pmatrix} 0 & 1 \\ -1 & s \end{pmatrix}\right) = -ud^2 - uc^2 + s(r - ucd).$$

By Lemma 2.16 (ii), we have the set $\{-ud^2 - uc^2 + s(r - ucd) \mid c, d \in \mathcal{F}, (c, d) \neq (0, 0)\}$ has $q - 1$ elements and thus (ii) follows. $\square$

Lemma 2.19. Let $A = \begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}$, $B = \begin{pmatrix} 0 & 1 \\ -1 & v \end{pmatrix}$, where $v, w \in \mathcal{F}$ are such that $v^2 - 4$ and $w^2 - 4$ are both non-square elements, i.e. A and B are of type (iv).

Given any f in $\{-i^2 + i(v - w) + w - 2 \mid i \in \mathcal{F}\}$, there exists a matrix E in the product $A^S B^S$ such that $\text{Trace}(E) = f$.

Let s be a non-square element of $\mathcal{F}$. If $v + w \neq 0$, the matrices $\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}$ and $\begin{pmatrix} -1 & s \\ 0 & -1 \end{pmatrix}$ are in the product $A^S B^S$. Otherwise the matrices $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $\begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix}$ are in the product $A^S B^S$.

We conclude that given any conjugacy classes A^S, B^S in $\mathcal{S}$, where both A and B are of type (iv), the product $A^S B^S$ is the union of at least $\frac{q+3}{2}$ distinct conjugacy classes. $\eta(A^S B^S) \geq \frac{q+3}{2}$.

Proof. Given any $i \in \mathcal{F}$, let $C(i) = \begin{pmatrix} 1 & i \\ 0 & 1 \end{pmatrix}$. Then by Lemma 2.5 (vi), we have that

$$\text{Trace}(A^{C(i)} B) = -i^2 + i(v - w) + w - 2.$$

Fix a and c in $\mathcal{F}$, where $a \neq 0$. Set $t_1 = -a^2 - c^2 + acw$ and $C = \begin{pmatrix} a & \frac{c-aw}{t_1} \\ c & -\frac{a}{t_1} \end{pmatrix}$.

We can check that $t_1 \neq 0$ since $w^2 - 4$ is a non-square and $C \in \mathcal{F}$. Also

$$A^C = \begin{pmatrix} 0 & 1 \\ -1 & w \end{pmatrix}^C = \begin{pmatrix} w & -\frac{1}{t_1} \\ t_1 & 0 \end{pmatrix}.$$

Fix e and g in $\mathcal{F}$, where $e \neq 0$. Set $t_2 = -e^2 - g^2 + egv$ and $D = \begin{pmatrix} e & \frac{g}{t_2} \\ g & \frac{-e+gv}{t_2} \end{pmatrix}$. By Lemma 2.16 (iii), for any $e, g \in \mathcal{F}$ with $e \neq 0$, $t_2 \neq 0$ since $v^2 - 4$ is a non-square.

$$B^D = \begin{pmatrix} 0 & 1 \\ -1 & v \end{pmatrix}^D = \begin{pmatrix} 0 & -\frac{1}{t_2} \\ t_2 & v \end{pmatrix}.$$

Thus

$$A^C B^D = \begin{pmatrix} -\frac{t_2}{t_1} & -\frac{w}{t_2} - \frac{v}{t_1} \\ 0 & -\frac{t_1}{t_2} \end{pmatrix}.$$

Therefore, if $t_1 = t_2$ we get that $A^C B^D = \begin{pmatrix} -1 & -\frac{w+v}{t_1} \\ 0 & -1 \end{pmatrix}$. By Lemma 2.16 (iii), we have that $\{a^2 + c^2 - acw \mid a, c \in \mathcal{F}, a \neq 0\} = \mathcal{F} \setminus \{0\}$. Thus the set $\{-\frac{w+v}{t_1} \mid t_1 = -a^2 - c^2 + acw, a, c \in \mathcal{F}, a \neq 0\}$ has $q - 1$ elements as long as $w + v \neq 0$. If $w + v = 0$, then $w - v \neq 0$ since $w \neq 0$. In that case, let $t_1 = -t_2$ and thus the set $\{-\frac{w-v}{t_1} \mid t_1 = -a^2 - c^2 + acw, a, c \in \mathcal{F}, a \neq 0\}$ has $q - 1$ elements. In particular, in both cases the sets contain 1 and a non-square element and the result follows. $\square$

Proof of Theorem B. If at least one of A and B is a scalar matrix, then $A^S B^S$ is a conjugacy class. We may assume then that A, B are similar to matrices of type (ii), (iii) or (iv). Theorem B then follows from Lemma 2.7, Lemma 2.18 and Lemma 2.19. $\square$

Proposition 2.20. *Assume that $\mathcal{F}$ is a field of q elements, with $q > 3$ odd.*

(i) *Assume that $q \cong 1 \pmod{4}$. Let w be a non-square element in $\mathcal{F}$. Let $A = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix}$. Then*

$$(2.21) \quad \eta(A^S B^S) = \frac{q+3}{2}.$$

(ii) *Assume that $q \not\cong 1 \pmod{4}$. Set $E = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$. Then*

$$(2.22) \quad \eta(E^S E^S) = \frac{q+3}{2}.$$

Hence, Theorem B is optimal.

Proof. (i) Let $C_i = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix} \in \mathcal{S}$, for $i = 1, 2$. By Lemma 2.4 (ii),

$$A^{C_i} B = \begin{pmatrix} 1 + c_i d_i & d_i^2 \\ -c_i^2 & 1 - c_i d_i \end{pmatrix} \begin{pmatrix} 1 & w \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 + c_i d_i & w + w c_i d_i + d_i^2 \\ -c_i^2 & 1 - c_i d_i - w c_i^2 \end{pmatrix}.$$

Hence, $\text{Trace}(A^{C_i} B) = 2 - w c_i^2$, which takes on $\frac{q+1}{2}$ values by Lemma 2.15. Note that if two matrices $A^{C_1} B$ and $A^{C_2} B$ have the same trace, then $c_1^2 = c_2^2$.

Suppose $c_1^2 = c_2^2 \neq 0$. Let $D = \begin{pmatrix} 1 & \frac{c_2 d_2 - c_1 d_1}{c_1^2} \\ 0 & 1 \end{pmatrix}$.

Then $(A^{C_1} B)^D = A^{C_2} B$, and so, excluding trace 2, each possible value of the trace is obtained by at most one conjugacy class.

Suppose $c_1 = c_2 = 0$. Then $A^{C_i}B = \begin{pmatrix} 1 & w + d_i^2 \\ 0 & 1 \end{pmatrix}$. Note that $w + d_i^2 \neq 0$ since $-w$ is non-square. Let z be a generator of the multiplicative group of units of $\mathcal{F}$, and suppose $w + d_i^2 = z^{n_i}$. By Lemma 2.14, $A^{C_1}B$ and $A^{C_2}B$ are conjugate if and only if there is an $e \in \mathcal{F} \setminus \{0\}$ such that $z^{n_1}e^2 = z^{n_2}$, i.e. when n_1 and n_2 have the same parity. Hence there are at most two conjugacy classes represented by matrices with trace 2, and thus, $\eta(A^S B^S) \leq (\frac{q+1}{2} - 1) + 2 = \frac{q+3}{2}$.

(ii) Define C_i as in (i). Then we may proceed by replacing w with 1 in the argument for the previous case, since -1 is not a square in $\mathcal{F}$. Thus, $\eta(E^S E^S) \leq \frac{q+3}{2}$.

Hence, in each case, the result follows by Theorem B. $\square$

REFERENCES

- [1] E. Adan-Bante, H. Verrill, Symmetric groups and conjugacy classes, J. Group Theory 11 (2008), no. 3, 371-379.
- [2] E. Adan-Bante, J. Harris, H. Verril, Products of conjugacy classes of the alternating group, preprint.
- [3] Z. Arad, M. Herzog, Products of conjugacy classes in groups, volume 1112 of Lecture notes in mathematics, Springer-Verlag, 1985.
- [4] G. D. James, M. Liebeck, Representations and Characters of Groups, Mathematics - 2001 - 466 pages.
- [5] The GAP Group, *GAP - Groups, Algorithms, and Programming, Version 4.4.10*; 2007, (<http://www.gap-system.org>).
- [6] R. Gow, Commutators in finite simple groups of Lie type, Bull. London Math. Soc. 32 (2000), 311-315.

DEPARTMENT OF MATHEMATICAL SCIENCE, NORTHERN ILLINOIS UNIVERSITY, WATSON HALL
320 DEKALB, IL 60115-2888, USA

E-mail address: EdithAdan@illinoisalumni.org

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN MISSISSIPPI, 730 EAST BEACH
BOULEVARD, LONG BEACH, MS 39560, USA

E-mail address: john.m.harris@usm.edu